

The planar Chain Rule and the Differential Equation for the planar Logarithms

L. Gerritzen

(29.09.2004)

Abstract

A planar monomial is by definition an isomorphism class of a finite, planar, reduced rooted tree. If x denotes the tree with a single vertex, any planar monomial is a non-associative product in x relative to m -array grafting. A planar power series $f(x)$ over a field K in x is an infinite sum of K -multiples of planar monomials including the unit monomial represented by the empty tree.

For every planar power series $f(x)$ there is a universal differential $df(x)$ which is a planar power series in x and a planar polynomial in a variable y which is the differential dx of x .

We state a planar chain rule and apply it to prove that the derivative $\frac{d}{dx}(Exp_k(x))$ is the k -ary planar exponential series. A special case of the planar chain rule is proved and it derived that the planar universe series $Log_k(1+x)$ of $Exp_k(x)$ satisfies the differential equation

$$\left((1+x) \frac{d}{dx} \right) (Log_k(1+x)) = 1$$

where $(1+x) \frac{d}{dx}$ is the derivative which when applied to x results in $1+x$.

Introduction

For every planar power series $f(x)$ in one variable x , there is a universal differential $df(x)$ which is a planar power series in x and a planar polynomial in a variable y which is the differential dx of x .

If $g(x)$ is a planar power series of order ≥ 1 , then the substitution of $g(x)$ for x in $f(x)$ gives a power series $f(g(x))$ and the differential $df(g(x))$ of $f(g(x))$ can be computed as

$$d\varphi_g(df)$$

where φ_g is the substitution homomorphism induced by g and $d\varphi_g$ is the homomorphism on the algebra of universal differential forms in x and dx extending φ_g

and mapping dx into dg .

This formula is called the planar chain rule.

We note an application to the planar exponential series $Exp_k(x)$, $k \in \mathbb{N}$, $k \geq 2$, see [G1], section 3.

One obtains a functional equation for the differential ω_k of $Exp_k(x)$, see section 6. It follows that the derivative $\frac{d}{dx}(Exp_k(x))$ is equal to $Exp_k(x)$.

This gives a conceptual proof of this differential relation, while the proof in [G1], section 3, is quite technical.

We also prove a special chain rule, if $g'(x)$ where $g'(x) = \frac{d}{dx} g(x)$ is the derivative with respect to x . From it we find that

$$\left((1+x) \frac{d}{dx} \right) (Log_k(1+x)) = 1,$$

if $Log_k(1+x)$ is the k -ary planar logarithm, see [G 2], section 5.

In section 1 notions about $\{x, y\}$ -labeled finite, planar, reduced rooted trees are fixed. Some basic facts about the algebra $K\{\{x, y\}\}$ of power series in x and polynomials in y over a field K are presented in section 2. The universal differential

$$d : K\{\{x\}\} \rightarrow K\{\{x, y\}\}$$

is introduced in section 4 and the planar chain rule is proved in section 5.

The application to the derivative of $Exp_k(x)$ relative to x is given in section 6.

In section 7 we prove a special rule for the derivative $\frac{d}{dx}$, if the substitution series $g(x)$ satisfies the equation $\frac{d}{dx} g(x) = 1 + g(x)$ and we show that

$$\left((1+x) \frac{d}{dx} \right) (Log_k(1+x)) = 1.$$

If h_n is the homogeneous component of $Log_k(1+x)$ of degree n , then

$$h'_{n+1}(x) = (-n)h_n(x)$$

for $n \geq 1$.

In [G2], section 6, the homogeneous component h_4 of $Log_k(1+x)$ has been computed to be

$$\begin{aligned} h_4(x) = & \left(\frac{k-3}{3![2]} - \frac{1}{4![3]}(k+1)(k-2)(k-3) \right) \cdot x^4 + \\ & + \left(\frac{1}{2} \frac{1}{3![2]} - \frac{2}{4![3]}(k-2) \right) (x \cdot x^3 + x^3 \cdot x) \end{aligned}$$

$$\begin{aligned}
& + \left(\frac{1}{3![2]!} \cdot \frac{3}{2} - \frac{1}{8} - \frac{3}{4![3]!} \right) (x \cdot (x \cdot x^2) + x \cdot (x^2 \cdot x) + (x \cdot x^2) \cdot x + (x^2 \cdot x) \cdot x) \\
& + \left(\frac{1}{3![2]!} \cdot \frac{3}{2} - \frac{1}{8} - \frac{3(k+1)}{4![3]!} \right) \cdot x^2 \cdot x^2 \\
& + \left(\frac{1}{2} \frac{k-2}{3![2]!} - \frac{2(k+1)(k-2)}{4![3]!} \right) (x \cdot x \cdot x^2 + x \cdot x^2 \cdot x + x^2 \cdot x \cdot x).
\end{aligned}$$

From this it follows that

$$\begin{aligned}
h_3(x) &= \left(-\frac{1}{3}\right)h'_4(x) = \frac{1}{4} \frac{k}{[2]} (x \cdot x^2 + x^2 \cdot x) - \frac{1}{3!} \frac{k-2}{[2]} \cdot x^3, \\
h_2(x) &= \left(-\frac{1}{2}\right)h'_3(x) = -\frac{1}{2} x^2, \\
h_1(x) &= -h'_2(x) = x.
\end{aligned}$$

1 Labeled planar rooted trees

Let x, y be different elements; they will be called variables in the sequel.

A $\{x, y\}$ -labeled finite, planar reduced rooted tree is a pair $S = (T, \lambda)$ where T is a finite, planar, reduced rooted tree, see [G1], and λ is a map $L(T) \rightarrow \{x, y\}$ where $L(T)$ denotes the set of leaves of T .

Then

$$\deg_x(T, \lambda) := \#\lambda^{-1}(x)$$

is called the x -degree of (T, λ) which is the number of leaves of T labeled with x by λ .

Also

$$\deg_y(T, \lambda) := \#\lambda^{-1}(y)$$

is called the y -degree of (T, λ) which is the number of leaves of T labeled with y by λ . Then $\deg(T, \lambda) = \deg T := \#L(T)$ is called the total degree of (T, λ) .

We denote by $P(x, y)$ the set of isomorphism classes of $\{x, y\}$ -labeled finite, planar, reduced rooted trees.

Given $S_1, \dots, S_m \in P(x, y)$, $S_i = (T_i, \lambda_i)$ for $1 \leq i \leq m$. Then there is a unique $S \in P(x, y)$ with the following properties:

$S = (T, \lambda)$ and if ρ_T is the root of T , then $T - \rho_T$ is a planar rooted forest whose i -th component is isomorphic with T_i . The set of leaves of $L(T)$ is the disjoint union of $L(T_1), \dots, L(T_n)$ and the labeling λ induces λ_i on T_i for all $1 \leq i \leq m$.

We recall that a planar rooted forest is an ordered disjoint system of planar rooted trees.

The construction described above gives a unique map

$$\bullet_m : (P(x, y))^m \rightarrow P(x, y)$$

which is called the m -ary grafting operation for labeled planar rooted trees.

Let $P'(x, y) = P(x, y) \cup \{1_P\}$ where 1_P is a symbol representing the empty tree.

There is a unique extension of the m -ary grafting operation to a map

$$\bullet_m : (P'(x, y))^m \rightarrow P'(x, y)$$

satisfying the properties: $\cdot_m(1_P, \dots, 1_P) = 1_P$

$\cdot_2(S, 1_P) = \cdot_2(1_P, S) = S$ for all $S \in P'(x, y)$,

$\cdot_m(S_1, \dots, S_m) = \cdot_{m-1}(S_1, \dots, S_{i-1}, S_{i+1}, \dots, S_m)$ if $S_i = 1_P$ for all $S_i \in P'(x, y)$.

It is also referred to as the m -ary grafting operation.

2 Polynomials and Power series

Let K be a field and V be the K -vectorspace of all K -valued functions f on $P'(x, y)$ such that for all $n \in \mathbb{N}$ the set

$$\{S \in P'(x, y) : f(S) \neq 0, \text{ord}_x(S) = n\}$$

is finite. For any $f \in V$ the value $f(S)$ at $S \in P'(x, y)$ will be denoted by $c_S(f)$. It will be called the coefficient of f relative to S .

For any $m \in \mathbb{N}, m \geq 2$, we are going to define a m -ary operation $\bullet_m$ on V by putting

$$f := \bullet_m(f_1, \dots, f_m)$$

if $c_S(f) := \sum c_{S_1}(f_1) \cdot \dots \cdot c_{S_m}(f_m)$ where the summation is extended over all $S_1, \dots, S_m$ such that $S = \bullet_m(S_1, \dots, S_m)$.

It is easy to check that $\bullet_m$ is a K -multilinear map $V^m \rightarrow V$. We will write $f_1 \times f_2 \times \dots \times f_m$ for $\bullet_m(f_1, \dots, f_m)$ sometimes.

In this note a K -algebra is by definition a K -vectorspace W together with a family $(\bullet_m)_{m \geq 2}$ of K -multilinear maps $\bullet_m : W^m \rightarrow W$. In the context of operad theory such an algebra can be defined as an algebra over an operad in the category of K -linear spaces which is freely generated by $\{\mu_m : m \geq 2\}$ where μ_m has degree m , see [MSS], Chap. (1.9). It is a tree operad, see [BV]. Trees are inspired by the attempt to obtain a general composite operation from a collection of indecomposable operations, see [BV].

Other authors, see [SV], use the term hyperalgebra to denote this type of algebras. For the general theory of operads and algebras over operads see also [L], [LR], [H], [F].

Therefore $(V, (\bullet_m)_{m \geq 2})$ is a K -algebra; it is denoted by $K\{\{x, y\}\}$ and will be referred to as the algebra of planar power series in x and polynomials in y . For any $S \in P'(x, y)$ there is a unique $f_S \in K\{\{x, y\}\}$ such that

$$c_T(f_s) = \begin{cases} 1 & : T = S \\ 0 & : T \in P'(x, y), T \neq S. \end{cases}$$

We identify f_S with S and thus consider $P'(x, y)$ as a subset of $K\{\{x, y\}\}$. It is easy to check that $P'(x, y)$ is a K -linearly independent subset of $K\{\{x, y\}\}$ and that the K -vectorspace of $K\{\{x, y\}\}$ generated by $P'(x, y)$ is a subalgebra. It is denoted by $K\{\{x, y\}\}$ and is called the algebra of planar polynomials: in x and y over K .

For $f \in K\{\{x, y\}\}$ define

$$\text{ord}_x(f) = \infty, \text{ if } f = 0$$

and if

$$f \neq 0,$$

then

$$\text{ord}_x(f) = \min\{n \in \mathbb{N} : \text{there is } S \in P(x, y) \text{ with } \deg_x(S) = n \text{ and } c_S(f) \neq 0.\}$$

It is called the x -order of f

Let

$$|f|_x := \begin{cases} 0 & : f = 0 \\ (\frac{1}{2})^{\text{ord}_x(f)} & : f \neq 0. \end{cases}$$

Then $|f_1 \cdot f_2 \cdot \dots \cdot f_m|_x = |f_1|_x \cdot |f_2|_x \cdot \dots \cdot |f_m|_x$

as $\text{ord}(f_1 \cdot f_2 \cdot \dots \cdot f_m) = \sum_{i=1}^m \text{ord}(f_i)$ and $|\cdot|_x$ defines a distance function on $K\{\{x, y\}\}$ by $|f, g|_x := |f - g|_x$ which is called the x -adic distance.

Proposition 2.1. *Any Cauchy sequence in $K\{\{x, y\}\}$ relative to the x -adic distance is convergent.*

Proof. By standard arguments. □

3 Substitution homomorphisms

Proposition 3.1. *Let $g, h \in K\{\{x, y\}\}$ and let $\text{ord}_x(g) \geq 1$. Then there is a unique K -algebra homomorphism*

$$\varphi_{(g, h)} = \varphi : K\{\{x, y\}\} \rightarrow K\{\{x, y\}\}$$

such that

$$(i) \quad \varphi(x) = g$$

$$(ii) \quad \varphi(y) = h$$

(iii) φ is continuous with respect to the x -adic topology.

It is called substitution homomorphism induced by (g, h) .

Proof. 1) First we define $\varphi'(S)$ for any $S \in P(x, y)$. If $S = (T, \lambda)$ where T is a planar rooted tree of degree m , there is a m -ary operation $\cdot_T$ associated to T , see [G2], Proposition (2.7)

Let $\varphi'(S) := \cdot_T(Z_1, \dots, Z_m)$ where $Z_i = g$ if $\lambda(l_i) = x$ and $Z_i = h$ if $\lambda(l_i) = y$. Here l_i is the i -th leaf of T .

2) There is a unique K -linear extension of the map φ' in 1) to a linear map

$$\varphi : K\{\{x, y\}\} \rightarrow K\{\{x, y\}\}.$$

3) If $f \in K\{\{x, y\}\}$, $f_n := \sum c_S(f) \cdot S$ where the sum is extended over all $S \in P(x, y)$ for which $\deg_x(S) = n$, then f_n is a finite sum and

$$f = \sum_{n=0}^{\infty} f_n.$$

From 1) it follows that $\text{ord}_x(\varphi(f)) \geq \text{ord}_x(f)$ for all $f \in K\{\{x, y\}\}$ as $\text{ord}_x(g) \geq 1$. Thus $\text{ord}_x(\varphi(f_n)) \geq n$ for all n .

One defines $\varphi(f) = \sum_{n=0}^{\infty} \varphi(f_n)$. The infinite sum is converging, as the sequence of partial sums is a Cauchy sequence with respect to $|\cdot|_x$.

Thus a continuous K -linear map $\varphi : K\{\{x, y\}\} \rightarrow K\{\{x, y\}\}$ is well defined. One can check that φ is an algebra homomorphism because its restriction to $K\{\{x, y\}\}$ is an algebra homomorphism by the definition in 1) and 2).

□

Notation: For $\varphi = \varphi(g, h)$ and $f \in K\{\{x, y\}\}$, we denote $\varphi(f)$ also by $f(g, h)$. Then $f = f(x, y)$ as $\varphi(x, y) = \text{id}$. Also $f(g, h) = f(g(x, y), h(x, y))$. It is obtained by replacing x by g and y by h .

For any $k \in K$ we get $f(kx, ky) = k^m f(x, y)$, if f is homogeneous of degree m .

4 Universal derivation

We denote by $K\{\{x\}\}$ the closed unital K -subalgebra of $K\{\{x, y\}\}$ generated by x .

Proposition 4.1. *There is a unique continuous K -linear map*

$$d : K\{\{x\}\} \rightarrow K\{\{x, y\}\}$$

such that

$$(i) \quad d(x) = y$$

$$(ii) \quad d(f_1 \cdot f_2 \cdot \dots \cdot f_m) = \sum_{i=1}^m (f_1 \cdot \dots \cdot f_{i-1} \cdot d(f_i) \cdot f_{i+1} \cdot \dots \cdot f_m) \text{ for all } m \geq 2 \text{ and all } f_1, \dots, f_m \in K\{\{x, y\}\}$$

The map d is called the universal derivation on $K\{\{x\}\}$. It is customary to denote $d(f)$ by df and call it the differential of f . Especially $y = dx$.

Proof. 1) First $d'(S)$ is defined for all $S \in P(x, y)$. If $S = 1_P$, then $d'(1_P) := 0$. If $S \neq 1_P$, $S = (T, \lambda)$, $\lambda(l_i) = x$ for all leaves $l_1, \dots, l_m$ of T , $m = \deg(T)$, $L(T) = \{l_1, \dots, l_m\}$, then

$$\text{let } d'(S) := \sum_{i=1}^m (T, \lambda^{(i)}), \text{ where } \lambda^{(i)}(l_j) = y \text{ if } j = i \text{ and } \lambda^{(i)}(l_j) = x \text{ if } j \neq i.$$

2) As $P'(x)$ is K -linearly independent, the map d' in 1) has a unique extension to a K -linear map

$$d_0 : K\{x\} \rightarrow K\{x, y\}.$$

3) There is a unique continuous extension of d_0 to a K -linear map

$$d : K\{\{x\}\} \rightarrow K\{\{x, y\}\}.$$

It is easy to check that property (ii) holds. □

5 Chain rule

Let $g \in K\{\{x\}\}$, $\text{ord}(g) \geq 1$. As a special case of Proposition (3.1) we get a continuous K -algebra homomorphism

$$\varphi_g = \varphi : K\{\{x\}\} \rightarrow K\{\{x\}\}$$

such that $\varphi(x) = g$.

Also there is a unique continuous K -algebra homomorphism

$$dy : K\{\{x, y\}\} \rightarrow K\{\{x, y\}\}$$

such that

$$d\varphi(x) = \varphi(x) = g$$

$$d\varphi(y) = dg$$

Proposition 5.1.

$$(d\varphi) \circ d = d \circ \varphi.$$

Proof. 1) Let $\delta := (d\varphi) \circ d - d \circ \varphi$. Then δ is a continuous K -linear map from $K\{\{x\}\}$ into $K\{\{x, y\}\}$ which satisfies

$$\delta(x) = 0$$

2) It is easy to see that

$$\delta(f_1 \cdot \dots \cdot f_m) = \sum_{i=1}^m \varphi(f_1) \cdot \dots \cdot \varphi(f_{i-1}) \cdot \delta(f_i) \cdot \varphi(f_{i+1}) \cdot \dots \cdot \varphi(f_m).$$

Using this formula one can prove by induction on the degree that for any $S \in P(x)$ one gets $\delta(S) = 0$.

If $S = S_1 \cdot \dots \cdot S_m$, then $\deg_x(S_i) < \deg_x(S)$ and by induction hypothesis $\delta(S_i) = 0, (\forall i?)$ By the formula above then $\delta(S) = 0$

3) As δ is continuous and $K\{x\}$ is dense in $K\{\{x\}\}$ and $\delta(f) = 0$ by 2) and the linearity of δ , we see that $\delta = 0$. $\square$

Example: Let

$$g = k \cdot x, k \in K.$$

Then

$$\varphi(f(x)) = f(kx)$$

and

$$(d\varphi)\omega(x, y) = \omega(kx, ky).$$

If f (resp. ω) is homogeneous of degree m , then $f(kx) = k^m f(x), \omega(kx, ky) = k^m \omega(x, y)$. $\square$

6 The derivative of the exponentials

Let now $\text{char}(K) = 0, k \in \mathbb{N}, k \geq 2$, and $f = \text{Exp}_k(x)$ be the k -ary exponential series, see [G1].

Let $\omega = df$ be the differential of f . Then $\omega(kx, ky) = df^k$ by the chain rule.

If $f = \sum_{n=0}^{\infty} f_n$ and f_n is the homogeneous part of degree n , then

$$\omega = \sum_{n=1}^{\infty} df_n$$

and df_n is the homogeneous component ω_n of ω of degree n with $df_0 = 0$.
More precisely:

$$\begin{aligned} \deg_x(df_n) &= n - 1, \\ \deg_y(df_n) &= 1. \end{aligned}$$

Now by the functional equation satisfied by f , we get

$$f^k(x) = f(kx) = \sum_{n=0}^{\infty} k^n f_n(x)$$

(*)

$$k^n f_n(x) = \sum_{i_1 + \dots + i_k = n} f_{i_1} \cdot f_{i_2} \cdot \dots \cdot f_{i_k}$$

where the summation is over all

$$(i_1, \dots, i_k) \in \mathbb{N}^k$$

with

$$i_1 + \dots + i_k = n.$$

Proposition 6.1.

$$k^n \omega_n(x, y) = \sum_{j=1}^k \sum_{i_1 + \dots + i_k = n} f_{i_1} \cdot \dots \cdot f_{i_{j-1}} \cdot \omega(f_{i_j}) \cdot f_{i_{j+1}} \cdot \dots \cdot f_{i_k}.$$

Proof. Immediate by property (ii) of Proposition 4.1. □

There is a unique continuous unital algebra homomorphism

$$\psi : K\{\{x, y\}\} \rightarrow K\{\{x\}\}$$

such that $\psi(x) = x, \psi(y) = 1$.

Recall that $\psi(f) = f$ for all $f \in K\{\{x\}\}$.

Then $(f \circ d) : K\{\{x\}\} \rightarrow K\{\{x\}\}$ is the

derivative relative to x , also denoted symbolically by $\frac{d}{dx}$, see [G1]

Proposition 6.2. $\frac{d}{dx}(Exp_k(x)) = Exp_k(x).$

Proof. 1) We have to show that the derivative $f'_n = \frac{d}{dx}(f_n)$ of the homogeneous component of $f(x) = Exp_k(x)$ is equal to f_{n-1} for $n \geq 1$.

By applying ψ to the formula in Proposition (6.1) we get

$$k^n f'_n(x) = \sum_{j=1}^k \sum_{i_1 + \dots + i_k = n} f_{i_1} \cdot \dots \cdot f_{i_{j-1}} \cdot f'_{i_j} \cdot f_{i_{j+1}} \cdot \dots \cdot f_{i_k}.$$

2) We will show that $f'_n(x) = f_{n-1}(x)$ for $n \geq 1$ by induction on n .

If $n = 1$, then it is obviously true as $\text{ord}_x(f - (1 + x)) \geq 2$.

Let now $n \geq 2$. As $f' i_j = 0$ if $i_j = 0$ and $f' i_j = f_{i_j-1}$ for $n > 0_j \geq 1$, we obtain

$$k^n f'_n(x) = \sum_{j=1}^k \sum_{\substack{i_1 + \dots + i_k = n \\ i_j \geq 1}} f_{i_1} \cdot \dots \cdot f_{i_{j-1}} \cdot f_{i_j-1} \cdot f_{i_{j+1}} \cdot \dots \cdot f_{i_k}$$

Now for any $1 \leq i \leq k$ we get $\sum_{\substack{i_1 + \dots + i_k = n \\ i_j \geq 1}} f_{i_1} \cdot \dots \cdot f_{i_{j-1}} \cdot f_{i_j} - 1 \cdot f_{i_{j+1}} \cdot \dots \cdot f_{i_k} =$

$$\sum_{i_1 + \dots + i_k = n-1} f_{i_1} \cdot f_{i_2} \cdot \dots \cdot f_{i_k} = k^{n-1} - f_{n-1}(x)$$

by (*)

It follows that

$$k^n f'_n(x) = k \cdot k^{n-1} \cdot f_{n-1}(x)$$

and

$$f'_n(x) = f_{n-1}(x).$$

□

7 Special chain rules and logarithm

Let $h \in K\{\{x\}\}$. Then there is a unique algebra homomorphism $\varphi_{(x,h)} = \varphi : K\{\{x, y\}\} \rightarrow K\{\{x\}\}$ such that $\varphi(x) = x$ and $\varphi(y) = h$. Then

$$\theta := \varphi \circ d : K\{\{x\}\} \rightarrow K\{\{x\}\}$$

is a derivation on $K\{\{x\}\}$ which means that is K -linear, continuous and satisfies the general product rule

$$\theta(f_1 \cdot \dots \cdot f_m) = \sum_{i=1}^m f_1 \cdot \dots \cdot f_{i-1} \cdot \theta(f_i) \cdot f_{i+1} \cdot \dots \cdot f_m$$

for all

$$f_1, \dots, f_m \in K\{\{x\}\}$$

This derivation θ will be formally denoted by $h \frac{d}{dx}$.

Be aware that $(h \frac{d}{dx})(f) \neq h \cdot \frac{d}{dx}(f)$ in general.

We denote $\frac{d}{dx}(f)$ also by f' and call it the derivative of f with respect to x .

Proposition 7.1. : Let $g \in K\{\{x\}\}$ such that $g' = 1 + g$

Then

$$\frac{d}{dx}(f(g(x))) = f'(g(x)) + (x \frac{d}{dx})(f)(g(x)) = ((1 + x) \frac{d}{dx})(f)(g(x))$$

Proof. :1) First we will prove the formula in case f is a tree T of degree n
If $n = 1$, then $T = x$ and $\frac{d}{dx}x = 1$ while $(1+x)\frac{d}{dx}(T) = 1+x$. Thus $f(g(x)) = g(x)$
and $\frac{d}{dx}(g) = 1 + g$ from which the formula follows. Let now $n > 1$. Then

$$\frac{d}{dx}(T) = \sum_{i=1}^n T^{(i)}$$

where $T^{(i)} = \bullet_T(z_1, \cdot, z_n)$ with

$$z_j = \begin{cases} x: j \neq i \\ 1_P: j = i \end{cases}$$

and

$$x \frac{d}{dx}(T) = n \times T$$

Also

$$\frac{d}{dx}(T(g(x))) = \sum_{i=1}^n S^i$$

with

$$S^{(i)} = \bullet_T(W_1, \dots, W_n)$$

and

$$w_j = \begin{cases} g(x): j \neq i \\ g'(x): j = i \end{cases}$$

As $g' = 1 + g$, we obtain from the multilinearity of $\bullet_T$, that

$$S(i) = \bullet_T(g, \dots, g) + \bullet_{T^{(i)}}(g, \dots, g)$$

From these computations it follows that the proposition holds, if f is a tree

2) From 1) one can extend the result to polynomials by multilinearity. As both sides of the formula are continuous in f the result follows for power series. $\square$

Let $\text{Log}_k(1+x)$ be the k -ary planar logarithm. see [G2], section 5.

Proposition 7.2. :

$$((1+x)\frac{d}{dx}(\text{Log}_k(1+x)) = 1$$

for all

$$k \in \mathbb{N}, \quad k > 2.$$

Proof. Let

$$g = \text{Exp}_k(x) - 1$$

Then

$$\text{Log}_k(\text{Exp}_k(x)) = x$$

From the special chain rule above, we can conclude that

$$(((1+x)\frac{d}{dx})(\text{Log}_k))\text{Exp}_k(x) = 1$$

From this the formula follows as $f(g(x)) = 1$ only if $f(x) = 1$. $\square$

Corollary 7.3. : Let h_n be the homogeneous component of $\text{Log}_k(1+x)$ of degree n .

Then

$$h_0 = 0$$

and

$$h'_{n+1} = -nh_n$$

for

$$n \geq 1$$

Proof. :

$$\text{Log}_k(1+x) = \sum_{n=0}^{\infty} h_n =: h$$

and

$$(1+x)\frac{d}{dx}(h_n) = h'_n + nh_n,$$

as

$$(s\frac{d}{dx})(h_n) = n \times h_n.$$

Moreover h'_n is homogeneous of degree $(n-1)$.

Thus

$$h'_{n+1} + nh_n = 0$$

for all

$$n \geq 1$$

because

$$(1+x)\frac{d}{dx}(h) = 1.$$

$\square$

Example 7.4. We give the homogeneous terms $h_n(x)$ of $\text{Log}_k(1+x)$ of degree $n \leq 4$.

Recall that

$$[n] = \frac{k^n - 1}{k - 1} = \sum_{i=0}^{n-1} k^i$$

and that

$$[n]! = \prod_{i=1}^n [i].$$

So especially

$$[2] = k + 1, [3] = k^2 + k + 1, [3]! = (k + 1)(k^2 + k + 1) = k^3 + 2k^2 + 2k + 1.$$

For any planar rooted tree T in P , let $\{T\}$ be the sum of all planar trees S in P whose underlying rooted tree is isomorphic to the underlying rooted tree of T .

One can show that $\text{Exp}_k(x)$ and $\text{Log}_k(x)$ are infinite sums over $\{T\}$ with $T \in P$. Then the homogeneous components $h_n(x)$ for $n \leq 4$ can be written as, see [G2]:

$$h_1(x) = x$$

$$h_2(x) = -\frac{1}{2} x^2$$

$$h_3(x) = \frac{1}{4} \frac{k}{[2]} \{x \cdot x^2\} - \frac{1}{3!} \frac{k-2}{[2]} \cdot x^3$$

$$h_4(x) = \left(\frac{k-3}{3![2]} - \frac{1}{4![3]}(k+1)(k-2)(k-3)\right) \cdot \{x^4\} +$$

$$+ \left(\frac{1}{2} \frac{1}{3![2]} - \frac{2}{4![3]}(k-2)\right) \{x \cdot x^3\}$$

$$+ \left(\frac{1}{3![2]} \cdot \frac{3}{2} - \frac{1}{8} - \frac{3}{4![3]}\right) \{x \cdot (x \cdot x^2)\}$$

$$+ \left(\frac{1}{3![2]} \cdot \frac{3}{2} - \frac{1}{8} - \frac{3(k+1)}{4![3]}\right) \cdot \{x^2 \cdot x^2\}$$

$$+ \left(\frac{1}{2} \frac{k-2}{3![2]} - \frac{2(k+1)(k-2)}{4![3]}\right) + \{x \cdot x \cdot x^2\}$$

Easy computations give

$$\frac{d}{dx}(\{x \cdot x \cdot x^2\}) = 3 \{x \cdot x^2\} + 6 \{x \cdot x \cdot x\}$$

$$\frac{d}{dx}(\{x^2 \cdot x^2\}) = 2 \{x \cdot x^2\}$$

$$\frac{d}{dx}(\{x \cdot (x \cdot x^2)\}) = 6 \{x \cdot x^2\}$$

$$\frac{d}{dx}\{x \cdot x^3\} = 2\{x^3\} + 3 \{x \cdot x^2\}$$

from which one can derive that

$$h'_4(x) = -3h_3(x)$$

Similarly one gets

$$h'_3(x) = -2h_2(x)$$

$$h'_2(x) = -h_1(x)$$

In [DG] $Log_2(x)$ was computed to be

$$Log_2(x) = x - \frac{1}{2}x^2 + \frac{1}{3}\left(\frac{1}{2}x \cdot x^2 + \frac{1}{2}x^2 \cdot x\right) -$$

$-\frac{1}{4}\left(\frac{4}{21}x \cdot (x \cdot x^2) + \frac{4}{21}x \cdot (x^2 \cdot x) + \frac{5}{21}x^2 \cdot x^2 + \frac{4}{21}(x \cdot x^2) \cdot x + \frac{4}{21}(x^2 \cdot x) \cdot x\right) +$
higher terms.

It is obtained by substituting 2 for k in $Log_k(x)$.

Open question: In there a procedure to construct $h_{n+1}(x)$ which is an integral of $(-n) \cdot h_n(x)$ directly from $h_n(x)$?

The problem arises from the fact that the space of homogeneous polynoms of degree $n + 1$ whose derivative is zero is non- trivial for $n \geq 2$

References

- [BV] Boardman, J. M. - Vogt, R. M.: Homotopy invariant algebraic structures on topological spaces, Springer-Verlag 1973
- [DG] Drensky, V.- Gerritzen, L.: Nonassociative exponential and logarithm, Journal of Algebra 272 (2004) 331 - 320
- [F] Fresse, B.: Koszul duality of operads and homology on partition posets, in "Homotopy theory and its applications (Evanston, 2002)", Contemp. Math. 346 (2004), 115-215.
- [G1] Gerritzen, L.: Planar rooted trees and non-associative exponential series, Advances in Applied Mathematics 33 (2004) 342 - 365
- [G2] Gerritzen, L.: Automorphisms of the planar tree power series algebra and the non-associative logarithm, Serdica Math. J., 30, Nos. 2 - 3 (2004), 135 - 158
- [Hol] Holtkamp, R.: On Hopf algebra structures over operads, preprint, July 2004, ArXiv:math.RA/0407074.

- [L] Loday, J. - L.: La renaissance des operades, Seninaire Burbaki, Vol. 1994/95, Asterisque No. 237 (1996) Exp. No. 792, 3, 47 - 74
- [LR3] Loday, J.-L. - Ronco, M.: Algebr de Hopf colibres, c. R. Acad. Sci. Paris, 337. (2003), 153-158.
- [MSS] Markl, M.- Shnider, S.- Stasheff, J.: Operads in Algebra, Topology and Physics, in: Math. Surveys Monogr., 2002, Chapter (1.5), p. 52 (see [G1]).
- [SU] Shestakov, F. P. - Umirbaev, U. U.: Free Akivis algebras, primitive elements and hyperalgebras, J. of Algebra 250 (2002), 533 - 548